



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the following named gentlemen were declared to be duly elected members of the Society :

Mr. Andrew Jackson Cassatt, of Philadelphia.

Mr. Clarence King, U. S. Geologist.

Mr. Horatio Hale, of Canada.

M. Paul Broca, M.D., of Paris.

Herr Franz Joseph Lauth, of Munich.

Dr. Isaac Norris, Jr., of Philadelphia.

And the meeting was adjourned.

ON A NEW VERTEBRATE GENUS FROM THE NORTHERN PART OF THE TERTIARY BASIN OF GREEN RIVER.

BY EDWARD D. COPE, A.M.

(Read before the American Philosophical Society, Oct. 18, 1872.)

ANAPTOMORPHUS ÆMULUS. Cope.

Dentition of the *ramus mandibuli*, In. 2, C. 1, P.M. 2, M. 3, total, 16; identical in number to those of *Simia* and *Homo*. It differs in many respects from these; there is no interruption in the series near the canine, and the symphysis though massive, is not co-ossified. Further details are, the last molar is three-lobed and elongated behind. The composition of the crowns of the preceding molars consists of four opposed lobes, which are very stout, and connected transversely by a thin ridge behind, or in close contact in front. The premolar tooth which is best preserved, is a perfect second, which, while having two roots, possesses a crown which stands almost entirely on the anterior, presenting a curved sectorial crest forwards and upwards.

<i>Measurements.</i>	<i>M.</i>
Length dental line.....	0.0148
“ of last molar.....	.0030
“ “ ante-penult.....	.0025
Width of “.....	.0020
Length of three molars preserved.....	.0070

DESCRIPTIONS OF NEW EXTINCT REPTILES FROM THE UPPER GREEN RIVER EOCENE BASIN, WYOMING.

BY E. D. COPE.

CROCODILUS (ICHTHYOSUCHUS) SUBULATUS. Cope, sp. nov.

Some of the cervical vertebræ without hypapophyses. Their cups round. Dentition peculiar. One or two very long smooth compressed straight teeth in the front of the *ramus mandibuli*. These are followed abruptly by a closely set series of sub-equal teeth of not one-fourth the